arXiv:1209.6104v1 [math.AP] 27 Sep 2012

FRACTIONAL LAPLACIAN ON THE TORUS

LUZ RONCAL AND PABLO RAÚL STINGA

Abstract. In this paper we study the fractional Laplacian $(-\Delta)^{\sigma/2}$ on the torus $\mathbb{T}$. Our approach is based on the semigroup language and the extension problem. We first obtain a pointwise integro-differential formula for $(-\Delta)^{\sigma/2}f(x)$, $0<\sigma<2$, $x\in\mathbb{T}$, that is given in terms of the classical Poisson integral on the unit disk. Sharp estimates for the kernel are given and the limits as $\sigma\to 0^+$ and $\sigma\to 2$ are computed exactly. Regularity estimates on Hölder, Lipschitz and Zygmund spaces are deduced. Secondly, we present a general extension problem to characterize any fractional power of a second order differential operator L^γ, $\gamma>0$. This generalizes previous known results in the case $0<\gamma<1$. The extension problem and the theory of degenerate elliptic equations is applied to prove interior and boundary Harnack's inequalities for $(-\Delta)^{\sigma/2}$.

1. Introduction

Very recently, there has been an increasing interest in the study of nonlinear partial differential equations involving fractional operators. The reason is that such problems arise naturally in applications, like Fluid Dynamics [5, 16], Strange Kinetics and Anomalous Transport [15], Financial Mathematics [3, 17], among many others. Fractional operators are very well-known from the Functional Analysis point of view, see the classical book by K. Yosida [22]. Nevertheless, there are some issues in these nonlinear fractional problems, not covered by the general theory, in which tools like pointwise formulas, Hölder estimates and Harnack's inequalities are needed [2, 3, 5, 16, 17].

In this paper we study fractional powers of the Laplacian $-\Delta=-\Delta_{\mathbb{T}}$ on the one-dimensional torus $\mathbb{T}$. Applying an approach based on the Poisson semigroup we obtain pointwise formulas for $(-\Delta)^{\sigma/2}f(x)$, $0<\sigma<2$, of the form

$$(-\Delta)^{\sigma/2}f(x)=\mathrm{P.V.}\int_{\mathbb{T}}(f(x)-f(y))K^{\sigma/2}(x-y)\,dy,$$

for a suitable singular kernel $K^{\sigma/2}(x)$. Moreover, by using these formulas, we show that

$$\lim_{\sigma\to 0^+}(-\Delta)^{\sigma/2}f(x)=f(x)-\frac{1}{2\pi}\int_{\mathbb{T}}f(y)\,dy,\qquad \lim_{\sigma\to 2}(-\Delta)^{\sigma/2}f(x)=-f''(x). \tag{1.1}$$

Observe the contrast of the first identity above with the case of the Laplacian $-\Delta_{\mathbb{R}^n}$ on $\mathbb{R}^n$, where $\lim_{\sigma\to 0^+}(-\Delta_{\mathbb{R}^n})^{\sigma/2}f(x)=f(x)$, see [19]. Taking into account the Fourier series definition of $(-\Delta)^{\sigma/2}f$ in (2.3) below, the identities in (1.1) are obvious, as limits in $L^2(\mathbb{T})$. But here we prove that the limits actually hold in the *pointwise* sense for a large class of functions. A crucial step for it is to compute all the constants exactly. We also study regularity properties of the fractional Laplacian on Hölder, Lipschitz and Zygmund spaces. On the other hand, by using the extension problem, we obtain interior and boundary Harnack's inequalities for $(-\Delta)^{\sigma/2}$.

The semigroup language we adopt here turns out to be the most adequate for our purposes. In particular, it allows us to compute all the constants exactly, to have an explicit solution for the extension problem and to study regularity properties in a simple and general way.

Date: First version: February 21, 2012.

2010 *Mathematics Subject Classification.* Primary: 35R11, 35B65, 35B45, 35R09. Secondary: 26A16, 47G20.

Key words and phrases. Fractional Laplacian, integro-differential operator, extension problem, degenerate elliptic equation, interior and boundary Harnack's inequalities, Hölder estimates, Poisson integral.

The first author was partially supported by grant MTM2009-12740-C03-03 from Ministerio de Ciencia e Innovación, Spain. The second author was partially supported by grant COLABORA 2010/01 from Planes Riojanos de I+D+I, Spain.

Apart from the results just mentioned in the concrete case of the fractional Laplacian on the torus, we also obtain the following generalization of the extension problems of [4] and [20]. Let L be a nonnegative self-adjoint second order partial differential operator with dense domain in $L^2(\Omega, d\eta)$, for some $\Omega \subseteq \mathbb{R}^n$ and a positive measure $d\eta$. Its fractional powers L^γ, $\gamma > 0$, can be defined through the spectral theorem.

Theorem 1.1. *Let $\gamma \in (0,\infty)\setminus\mathbb{N}$, and f in the domain of L^γ. A solution $u \in C^\infty((0,\infty); \mathrm{Dom}(L)) \cap C([0,\infty); L^2(\Omega))$ of the extension problem*

$$\begin{cases} -L_x u + \frac{1-2\gamma}{y} u_y + u_{yy} = 0, & \text{in } \Omega \times (0,\infty), \\ u(x,0) = f(x), & \text{on } \Omega, \end{cases} \tag{1.2}$$

is given by

$$u(x,y) = \frac{y^{2\gamma}}{4^\gamma \Gamma(\gamma)} \int_0^\infty e^{-tL} f(x)\, e^{-\frac{y^2}{4t}} \frac{dt}{t^{1+\gamma}}, \tag{1.3}$$

and

$$\lim_{y\to 0^+} y^{1-2(\gamma-[\gamma])} \partial_y \left((y^{-1}\partial_y)^{[\gamma]} u(x,y) \right) = C_\gamma L^\gamma f(x), \tag{1.4}$$

where $[\gamma]$ is the integer part of γ and

$$C_\gamma = \frac{4^{\gamma-[\gamma]}\Gamma(\gamma-[\gamma])}{2(\gamma-[\gamma])\Gamma(-(\gamma-[\gamma])} \cdot \frac{1}{2^{[\gamma]}(\gamma-[\gamma])(\gamma-[\gamma]+1)\cdots(\gamma-1)}.$$

The idea of introducing the semigroup language to study fine properties of fractional operators is contained in [20], where the Caffarelli-Silvestre extension problem is generalized to apply to general fractional operators L^γ, $0 < \gamma < 1$. For this range of γ, Theorem 1.1 above is already stated in [20]. The advantage of characterizing the nonlocal operator L^γ via (1.2) is that the extension equation is a local PDE for which well-known techniques from the Calculus of Variations can be applied, see [4]. Moreover, the part in y in the first equation of (1.2) is a Bessel operator, see [20].

Our novelty here is that γ can be taken to be any noninteger positive number. In this way we answer a question raised by Ricardo G. Durán about how to characterize higher-order fractional operators via the extension problem. When $L = -\Delta_{\mathbb{R}^n}$ and $0 < \gamma < 1$ we recover in (1.2) and (1.4) the extension problem of L. Caffarelli and L. Silvestre [4]. Also in this case, but for general noninteger $\gamma > 0$, condition (1.4) was obtained by S.-Y. A. Chang and M. d. M. González in [7]. They derive the result by relating the fractional Laplacian $(-\Delta_{\mathbb{R}^n})^\gamma$ with a class of conformally covariant operators P_γ defined at the boundary of the hyperbolic space. In turn, our approach here is more general and it is based on the explicit formula for u of (1.3), already found in [20] when $0 < \gamma < 1$, that involves the semigroup e^{-tL}.

In [4], Caffarelli and Silvestre used the extension problem and local PDE techniques to prove interior and boundary Harnack's estimates for $(-\Delta_{\mathbb{R}^n})^{\sigma/2}$. Since in this case the extension equation is a degenerate elliptic equation in $\mathbb{R}^{n+1}_+$, the theory by E. B. Fabes, D. Jerison, C. Kenig and R. Serapioni [9, 10] can be applied. Schauder and regularity estimates had been previously derived by Silvestre in [17]. These results were used to solve nonlinear problems for the fractional Laplacian, see for example [2, 3, 16, 17]. The extension problem technique was used in [20] to prove interior Harnack's inequality for the fractional harmonic oscillator $(-\Delta_{\mathbb{R}^n} + |x|^2)^{\sigma/2}$. In such case the extension equation becomes a degenerate elliptic Schrödinger equation, so Harnack's inequality proved by C. E. Gutiérrez in [11] was needed. In [21] the corresponding Hölder estimates for the fractional harmonic oscillator can be found.

The paper is divided into two parts. In the first one we focus on the definition of the fractional Laplacian on the torus, the pointwise formulas, the limits (1.1) and the interaction with Hölder, Lipschitz and Zygmund spaces. Our approach is based on the semigroup language. Up to our knowledge, there is not a closed formula for the heat semigroup on $\mathbb{T}$, so we work with the Poisson semigroup. The regularity estimates are also obtained with this general approach. Part 2 deals with the general extension problem and its applications. We prove Theorem 1.1 and we use it to find interior and boundary Harnack's inequalities for $(-\Delta)^{\sigma/2}$. The ideas therein are based on [4]. We point out here that interior Harnack's inequality for the fractional Laplacian on $\mathbb{R}^n$ is classical [12]. Boundary Harnack's estimates were first proved by K. Bogdan in [1] by using probabilistic techniques.

Throughout this paper the letters c and C denote positive constants that may change at each occurrence. For two positive quantities A and B we write $A \sim B$ to denote that there exist constants $0 < c \le C$ independent of σ such that $c \le A/B \le C$.

Part 1. Pointwise formulas and Hölder spaces

2. The fractional Laplacian on the torus

We denote by $\mathbb{T} \subset \mathbb{R}^2$ the unit circle. We will sometimes identify the torus with the interval $(-\pi, \pi]$ and functions on $\mathbb{T}$ with 2π-periodic functions on $\mathbb{R}$. Let Δ be the Laplace–Beltrami operator on the torus. It is well-known that $\Delta = \frac{d^2}{d\theta^2}$, where θ is the angular variable in polar coordinates $X = r\theta$, $r > 0$, $-\pi < \theta \le \pi$, in $\mathbb{R}^2$. Given an integrable function f on the torus $\mathbb{T}$, we write its Fourier series expansion as

$$f(x) = \sum_{n\in\mathbb{Z}} c_n(f) e^{inx}, \quad c_n(f) = \frac{1}{2\pi}\int_{\mathbb{T}} f(x) e^{-inx}\,dx.$$

The Poisson semigroup generated by $-\Delta$ is defined by

$$P_t f(x) \equiv e^{-t\sqrt{-\Delta}} f(x) := \sum_{n\in\mathbb{Z}} e^{-t|n|} c_n(f) e^{inx}, \quad f \in L^2(\mathbb{T}),\ t \ge 0.$$

Then $P_t f(x)$ is the solution of the harmonic extension $(\partial_{tt} + \Delta)v = 0$ in $\mathbb{T} \times (0,\infty)$, with boundary condition $v(x,0) = f(x)$ on $\mathbb{T}$. We have the convolution formula

$$P_t f(x) = \int_{\mathbb{T}} P_t(x-y) f(y)\,dy, \tag{2.1}$$

where

$$\begin{aligned} P_t(x) &= \frac{1}{2\pi}\sum_{n\in\mathbb{Z}} e^{-t|n|} e^{inx} = \frac{1}{\pi}\left(\frac{1}{2} + \sum_{n=1}^{\infty} e^{-tn}\cos nx\right) \\ &= \frac{1-e^{-2t}}{2\pi(1 - 2e^{-t}\cos x + e^{-2t})} = \frac{1-e^{-2t}}{2\pi((1-e^{-t})^2 + 4e^{-t}\sin^2(x/2))}, \end{aligned} \tag{2.2}$$

is the Poisson kernel. See Zygmund [23, Chapter III, (6.2)].

Let $f \in C^\infty(\mathbb{T})$. For $0 < \sigma < 2$, we define the fractional powers of the Laplacian on $\mathbb{T}$ as

$$(-\Delta)^{\sigma/2} f(x) = \sum_{n\in\mathbb{Z}} |n|^\sigma c_n(f) e^{inx}. \tag{2.3}$$

Note that for any $N \in \mathbb{N}$ there exists a constant C depending only on f such that $|c_n(f)| \le C|n|^{-N}$, for all n. Therefore, the series that defines $(-\Delta)^{\sigma/2} f$ is absolutely convergent and is a $C^\infty(\mathbb{T})$-function. We also have the symmetry property $\langle (-\Delta)^{\sigma/2} f, g\rangle_{L^2(\mathbb{T})} = \langle f, (-\Delta)^{\sigma/2} g\rangle_{L^2(\mathbb{T})}$, $g \in C^\infty(\mathbb{T})$. In fact, the series in (2.3) converges in $L^2(\mathbb{T})$ whenever f has the property that $\sum_{n\in\mathbb{Z}} |n|^{2\sigma} |c_n(f)|^2 < \infty$, that is, when f is in the Sobolev space $H^\sigma = \mathrm{Dom}((-\Delta)^\sigma)$. This allows us to extend the definition of $(-\Delta)^{\sigma/2}$ to this class. The sum in (2.3) is over $\mathbb{Z} \setminus \{0\}$ and the limits in (1.1) hold in $L^2(\mathbb{T})$.

It is easy to check that for any $\lambda > 0$ and $0 < \sigma < 2$ we have the integral identity

$$\lambda^{\sigma/2} = \frac{1}{c_\sigma}\int_0^\infty \left(e^{-t\lambda^{1/2}} - 1\right)^{[\sigma]+1} \frac{dt}{t^{1+\sigma}},$$

where

$$c_\sigma = \int_0^\infty (e^{-s} - 1)^{[\sigma]+1} \frac{ds}{s^{1+\sigma}}. \tag{2.4}$$

Plugging this into (2.3) and interchanging the summation with the integration, we get

$$(-\Delta)^{\sigma/2}f(x)=\frac{1}{c_\sigma}\int_0^\infty\left(e^{-t\sqrt{-\Delta}}-\mathrm{I}\right)^{[\sigma]+1}f(x)\,\frac{dt}{t^{1+\sigma}},\quad x\in\mathbb{T},\tag{2.5}$$

for $f\in C^\infty(\mathbb{T})$. Here I is the identity operator. It is not difficult to prove that the constant c_σ in (2.4) satisfies the following asymptotic estimates:

$$\frac{1}{c_\sigma}\sim(2-\sigma),\quad\text{as }\sigma\to2^-,\qquad\frac{1}{c_\sigma}\sim-\sigma,\quad\text{as }\sigma\to0^+.\tag{2.6}$$

3. Pointwise formulas for the fractional Laplacian

In this section we obtain explicit formulas for $(-\Delta)^{\sigma/2}f(x)$ when f belongs to the Hölder spaces. We also prove the pointwise limits in (1.1).

Definition 3.1 (Hölder spaces on $\mathbb{T}$)**.** Let $0<\alpha\le1$ and $k\in\mathbb{N}_0$. A continuous function $f:\mathbb{T}\to\mathbb{R}$ belongs to the Hölder space $C^{k,\alpha}(\mathbb{T})$, if $f\in C^k(\mathbb{T})$ and

$$[f^{(k)}]_{C^\alpha}:=\sup_{\substack{x,y\in\mathbb{T}\\x\neq y}}\frac{|f^{(k)}(x)-f^{(k)}(y)|}{\left|\sin\frac{x-y}{2}\right|^\alpha}<\infty.$$

We define the norm in the spaces $C^{k,\alpha}(\mathbb{T})$ to be $\|f\|_{C^{k,\alpha}(\mathbb{T})}:=\sum_{0\le l\le k}\|f^{(l)}\|_{L^\infty(\mathbb{T})}+[f^{(k)}]_{C^\alpha}$.

Remark 3.2. The condition $f\in C^{k,\alpha}(\mathbb{T})$ is equivalent to ask for f to be a 2π-periodic C^k function on $\mathbb{R}$ such that

$$|f^{(k)}(x)-f^{(k)}(y)|\le C_{k,\alpha}\left|x-y\right|^\alpha,$$

for all $x,y\in[-\pi,\pi]$ or, equivalently, for all $x,y\in\mathbb{R}$, $x\neq y$. The least constant $C_{k,\alpha}$ for which the inequality above holds is equivalent to $[f^{(k)}]_{C^\alpha}$.

Consider the test space $C^\infty(\mathbb{T})$ endowed with the family of norms

$$\|\phi\|_k^2:=\|(\mathrm{I}-\Delta)^k\phi\|_{L^2(\mathbb{T})}^2=\sum_{n\in\mathbb{Z}}(1+|n|^2)^k|c_n(\phi)|^2,\quad k\ge1,$$

A real linear functional S on $C^\infty(\mathbb{T})$ is a periodic distribution if it satisfies the following continuity property: if $\phi_j\in C^\infty(\mathbb{T})$, $\|\phi_j\|_k\to0$ as $j\to\infty$ for every $k\in\mathbb{N}$, then $S(\phi_j)\to0$. Note that if $f\in L^1(\mathbb{T})$ then f defines a periodic distribution by $f(\phi)=\int_{\mathbb{T}}f\phi$. See Schwartz [14, Chapter VII].

Observe that the fractional Laplacian is a continuous linear operator on $C^\infty(\mathbb{T})$. The following result will allow us to extend the definition of the fractional Laplacian to the Hölder classes.

Lemma 3.3. *Let T be a continuous linear operator on $C^\infty(\mathbb{T})$ such that $\langle T\phi,\psi\rangle_{L^2(\mathbb{T})}=\langle\phi,T\psi\rangle_{L^2(\mathbb{T})}$ for all $\phi,\psi\in C^\infty(\mathbb{T})$ and*

$$T\phi(x)=\int_{\mathbb{T}}(\phi(x)-\phi(y))K(x-y)\,dy,\quad\phi\in C^\infty(\mathbb{T}),\ x\in\mathbb{T}.$$

Assume that the kernel K above is a 2π-periodic function on $\mathbb{R}$ with

$$|K(x)|\le\frac{C_\gamma}{|x|^{1+\gamma}},\quad x\in\mathbb{T},\tag{3.1}$$

for some $0\le\gamma<1$. Let $f\in C^{0,\gamma+\varepsilon}(\mathbb{T})$, with $0<\gamma+\varepsilon\le1$, $\varepsilon>0$. Then Tf is well defined as a periodic distribution and it coincides with the continuous function

$$Tf(x)=\int_{\mathbb{T}}(f(x)-f(y))K(x-y)\,dy,\quad x\in\mathbb{T}.\tag{3.2}$$

Proof. By (3.1) and the assumption on f, the integral in (3.2) is absolutely convergent. As $f\in L^1(\mathbb{T})$, we can define Tf as a periodic distribution by using the symmetry of T via $(Tf)(\phi):=f(T\phi)=\int_{\mathbb{T}}fT\phi$, $\phi\in C^\infty(\mathbb{T})$. Let $f_j(x)=(P_{1/j}*f)(x)$, $j\in\mathbb{N}$, $x\in\mathbb{T}$, where $P_t(z)$ is the Poisson kernel (2.2). It is well known [23, Chapters III and IV] that $f_j\in C^\infty(\mathbb{T})$ and that $f_j\to f$, $j\to\infty$, in $L^p(\mathbb{T})$, $1\le p\le\infty$. It is not difficult to check that $[f_j]_{C^{\gamma+\varepsilon}}\le[f]_{C^{\gamma+\varepsilon}}$ for all j. Now, from the

$L^p(\mathbb{T})$-convergence of f_j to f, we can see that $Tf_j \to Tf$ as periodic distributions, that is to say $\lim_{j\to\infty}(Tf_j)(\phi) = \lim_{j\to\infty}\int_{\mathbb{T}} f_j T\phi = \int_{\mathbb{T}} f(T\phi) = (Tf)(\phi)$, for each $\phi \in C^\infty(\mathbb{T})$. Let $\eta > 0$. There exists $\delta > 0$ such that,

$$C_\gamma [f]_{C^{\gamma+\varepsilon}} \int_{|x-y|<\delta, |y|<\pi} |x-y|^{\varepsilon-1}\, dy < \frac{\eta}{3}.$$

Then, for all j,

$$\left| \int_{|x-y|<\delta} (f_j(x) - f_j(y))K(x-y)\, dy \right| + \left| \int_{|x-y|<\delta} (f(x) - f(y))K(x-y)\, dy \right| < \frac{2}{3}\eta.$$

On the other hand,

$$\left| \int_{\delta<|x-y|<\pi} \left[(f_j(x) - f_j(y)) - (f(x) - f(y))\right] K(x-y)\, dy \right|$$
$$\leq C\,|f_j(x) - f(x)| + C \left(\int_{\mathbb{T}} |f_j(y) - f(y)|^2\, dy \right)^{1/2} \leq \frac{\eta}{3},$$

for all sufficiently large j, uniformly in x in some compact subset of $\mathbb{T}$. Therefore the right hand side of (3.2) with f_j converges uniformly on compact subsets of $\mathbb{T}$ to the right hand side of (3.2) with f and the limit is a continuous function. By uniqueness of the limits, (3.2) holds. □

Next we derive the pointwise formulas for $(-\Delta)^{\sigma/2} f(x)$, see (3.3) and (3.9) below. In both cases the kernels are (3.4) and (3.10), respectively. Formally, from (2.5), it is evident that those are the correct kernels. The point is that (3.3) and (3.9) are valid for a large class of continuous functions and in the pointwise sense. As a consequence of this, the pointwise semigroup formula (2.5) is also true for Hölder continuous functions, see Remark 3.9 below.

Let us begin with the case $0 < \sigma < 1$.

Theorem 3.4 (Pointwise formula for $0 < \sigma < 1$)**.** *Let* $0 < \sigma < 1$ *and* $f \in C^{0,\sigma+\varepsilon}(\mathbb{T})$, *for some* $\varepsilon > 0$ *such that* $0 < \sigma + \varepsilon \leq 1$. *Then* $(-\Delta)^{\sigma/2} f$ *is a continuous function and*

$$(-\Delta)^{\sigma/2} f(x) = \int_{\mathbb{T}} (f(x) - f(y)) K^{\sigma/2}(x-y)\, dy, \tag{3.3}$$

where

$$K^{\sigma/2}(x) = \frac{1}{-c_\sigma} \int_0^\infty P_t(x) \frac{dt}{t^{1+\sigma}} \geq 0, \tag{3.4}$$

with c_σ *as in* (2.4). *Moreover, there exist universal constants* $C, c > 0$ *such that*

$$\frac{c}{-c_\sigma (1-\sigma) |\sin x/2|^{1+\sigma}} \leq K^{\sigma/2}(x) \leq \frac{C}{-c_\sigma \sigma (1-\sigma) |\sin x/2|^{1+\sigma}}. \tag{3.5}$$

Besides, the integral in (3.3) *is absolutely convergent.*

Proof. First we prove (3.5). Recall (2.2). Let us split the integral in (3.4) into two parts

$$J := \int_0^\infty \frac{1 - e^{-2t}}{2\pi((1-e^{-t})^2 + 4e^{-t}\sin^2(x/2))} \frac{dt}{t^{1+\sigma}} = \int_0^1 + \int_1^\infty =: J_1 + J_2.$$

For J_1,

$$J_1 \sim \int_0^1 \frac{t}{t^2 + C\sin^2(x/2)} \frac{dt}{t^{1+\sigma}} = \frac{1}{\sin^2(x/2)} \int_0^1 \frac{t^{-\sigma}}{\frac{t^2}{\sin^2(x/2)} + C}\, dt$$
$$= \frac{1}{|\sin(x/2)|^{1+\sigma}} \int_0^{1/|\sin(x/2)|} \frac{s^{-\sigma}}{s^2 + C}\, ds.$$

There exist constants $c, C > 0$ independent of σ such that

$$\int_0^{1/|\sin(x/2)|} \frac{s^{-\sigma}}{s^2 + C}\, ds \leq \frac{1}{C} \int_0^1 s^{-\sigma}\, ds + \int_1^\infty \frac{ds}{s^2 + C} \leq \frac{C}{1-\sigma},$$

and

$$\int_0^{1/|\sin(x/2)|} \frac{s^{-\sigma}}{s^2+C}\,ds \geq \frac{1}{1+C}\int_0^1 s^{-\sigma}\,ds = \frac{c}{1-\sigma}.$$

Thus $J_1 \sim \dfrac{1}{(1-\sigma)|\sin(x/2)|^{1+\sigma}}$. We also have $J_2 \leq C\displaystyle\int_1^\infty \frac{dt}{t^{1+\sigma}} = \frac{C}{\sigma}$. Therefore (3.5) follows.

To prove (3.3), suppose for the moment that $f \in C^\infty(\mathbb{T})$. Since

$$P_t 1(x) = \int_{\mathbb{T}} P_t(x-y)\,dy \equiv 1, \quad \text{for all } x \in \mathbb{T},\ t>0, \tag{3.6}$$

we have, by the formula with the Poisson semigroup in (2.5),

$$(-\Delta)^{\sigma/2} f(x) = \frac{1}{-c_\sigma}\int_0^\infty \int_{\mathbb{T}} P_t(x-y)(f(x)-f(y))\,dy\,\frac{dt}{t^{1+\sigma}}, \quad f \in C^\infty(\mathbb{T}). \tag{3.7}$$

Since $f \in C^\infty(\mathbb{T})$, by (3.5),

$$\int_0^\infty \int_{\mathbb{T}} |P_t(x-y)(f(x)-f(y))|\,dy\,\frac{dt}{t^{1+\sigma}} \leq C\int_{-\pi}^{\pi} \frac{|\sin\frac{z}{2}|}{|\sin\frac{z}{2}|^{1+\sigma}}\,dz < \infty.$$

Hence we can apply Fubini's theorem in (3.7) to obtain (3.3) for $f \in C^\infty(\mathbb{T})$. By Lemma 3.3, the same formula holds for $f \in C^{0,\sigma+\varepsilon}(\mathbb{T})$. □

We now consider the first pointwise limit in (1.1).

Proposition 3.5. *Let $f \in C^{0,\alpha}(\mathbb{T})$, for some $0<\alpha\leq 1$. Then*

$$\lim_{\sigma\to 0^+} (-\Delta)^{\sigma/2} f(x) = f(x) - \frac{1}{2\pi}\int_{\mathbb{T}} f(y)\,dy, \quad x \in \mathbb{T}.$$

Proof. We must check that

$$(-\Delta)^{\sigma/2} f(x) - f(x) + \frac{1}{2\pi}\int_{\mathbb{T}} f(y)\,dy = \int_{\mathbb{T}} (f(x)-f(y))\left[K^{\sigma/2}(x-y) - \frac{1}{2\pi}\right]dy \to 0,$$

as $\sigma \to 0^+$. Take any $0<\sigma<\alpha$. By (3.4),

$$\begin{aligned} K^{\sigma/2}(x) - \frac{1}{2\pi} &= \frac{1}{-c_\sigma}\int_0^1 P_t(x)\,\frac{dt}{t^{1+\sigma}} + \frac{1}{-c_\sigma}\int_1^\infty P_t(x)\,\frac{dt}{t^{1+\sigma}} - \frac{1}{2\pi} \\ &= \frac{1}{-c_\sigma}\int_0^1 P_t(x)\,\frac{dt}{t^{1+\sigma}} + \frac{1}{-c_\sigma}\int_1^\infty \left(P_t(x) - \frac{1}{2\pi}\right)\frac{dt}{t^{1+\sigma}} + \frac{1}{2\pi}\left(\frac{1}{-c_\sigma\sigma} - 1\right). \end{aligned} \tag{3.8}$$

As we did in the proof of Theorem 3.4, we can see that the first integral in (3.8) is bounded from above by $C\left(-c_\sigma(1-\sigma)|\sin(x/2)|^{1+\sigma}\right)^{-1}$. The third summand of (3.8) tends to zero as $\sigma \to 0^+$ because $-c_\sigma\sigma = -\sigma\Gamma(-\sigma) = \Gamma(1-\sigma)$. As for the second summand in (3.8), note that

$$\left|P_t(x) - \frac{1}{2\pi}\right| \leq Ce^{-t}\frac{1-e^{-t}}{(1-e^{-t})^2 + 4e^{-t}\sin^2(x/2)}.$$

Hence,

$$\frac{1}{-c_\sigma}\int_1^\infty \left|P_t(x) - \frac{1}{2\pi}\right|\frac{dt}{t^{1+\sigma}} \leq \frac{C}{-c_\sigma}\int_1^\infty e^{-t}\,dt = \frac{C}{-c_\sigma}.$$

Collecting terms in (3.8),

$$\left|K^{\sigma/2}(x) - \frac{1}{2\pi}\right| \leq \frac{C}{-c_\sigma(1-\sigma)|\sin(x/2)|^{1+\sigma}} + F(\sigma),$$

where $F(\sigma)$ is a function of σ that tends to zero as $\sigma \to 0^+$. Therefore,

$$\int_{\mathbb{T}} |f(x)-f(y)|\left|K^{\sigma/2}(x-y) - \frac{1}{2\pi}\right|dy \leq \frac{C}{-c_\sigma(1-\sigma)}\int_{\mathbb{T}} \frac{|f(x)-f(y)|}{|\sin\frac{x-y}{2}|^{1+\sigma}}\,dy + \|f\|_{L^\infty(\mathbb{T})}F(\sigma).$$

The second term above tends to zero as $\sigma \to 0^+$. For the first term, by applying (2.6),

$$\begin{aligned}\frac{C}{-c_\sigma(1-\sigma)}\int_{\mathbb{T}}\frac{|f(x)-f(y)|}{|\sin\frac{x-y}{2}|^{1+\sigma}}\,dy &\le \frac{C}{-c_\sigma(1-\sigma)}\int_{\mathbb{T}}\left|\sin\tfrac{x-y}{2}\right|^{\alpha-1-\sigma}dy\\ &\le \frac{C}{-c_\sigma(1-\sigma)} \to 0, \quad \text{as } \sigma\to 0^+.\end{aligned}$$

□

Next we take $1 \le \sigma < 2$.

Theorem 3.6 (Pointwise formula for $1 \le \sigma < 2$)**.** *Let $1 \le \sigma < 2$ and $f \in C^{1,\sigma+\varepsilon-1}(\mathbb{T})$, for some $\varepsilon > 0$ such that $0 < \sigma+\varepsilon-1 \le 1$. Then*

$$\begin{aligned}(-\Delta)^{\sigma/2}f(x) &= \int_{\mathbb{T}}(f(x)-f(y)-f'(x)(x-y))K^{\sigma/2}(x-y)\,dy\\ &= \mathrm{P.\,V.}\int_{\mathbb{T}}(f(x)-f(y))K^{\sigma/2}(x-y)\,dy \qquad (3.9)\\ &= \lim_{\delta\to 0}\int_{\delta\le\left|\frac{x-y}{2}\right|<\pi-\delta,|y|<\pi}(f(x)-f(y))K^{\sigma/2}(x-y)\ dy,\end{aligned}$$

where

$$K^{\sigma/2}(x) = \frac{1}{c_\sigma}\int_0^\infty (2P_t(x)-P_{2t}(x))\,\frac{dt}{t^{1+\sigma}} \ge 0, \tag{3.10}$$

and c_σ is as in (2.4). Moreover, there exist universal constants $C, c > 0$ such that

$$\frac{c}{c_\sigma|\sin x/2|^{1+\sigma}} \le K^{\sigma/2}(x) \le \frac{C}{c_\sigma|\sin x/2|^{1+\sigma}}. \tag{3.11}$$

Proof. Let us begin with the proof of (3.11). Let $D_t(x) := 2P_t(x)-P_{2t}(x)$. Then, by (2.2),

$$D_t(x) = \frac{(1-e^{-2t})(1-e^{-t})^2\left[2(1+e^{-t})^2-(1+e^{-2t})-4e^{-t}|\sin x/2|^2\right]}{2\pi((1-e^{-t})^2+4e^{-t}|\sin x/2|^2)((1-e^{-2t})^2+4e^{-2t}|\sin x/2|^2)} \ge 0.$$

We split the integral in (3.10) as $\int_0^1 + \int_1^\infty$. On one hand,

$$\begin{aligned}\int_0^1 D_t(x)\,\frac{dt}{t^{1+\sigma}} &\sim \int_0^1 \frac{t^3}{t^4+|\sin(x/2)|^4}\,\frac{dt}{t^{1+\sigma}}\\ &= \frac{1}{|\sin x/2|^{1+\sigma}}\int_0^{1/|\sin x/2|}\frac{s^{2-\sigma}}{s^4+1}\,ds \sim \frac{1}{|\sin(x/2)|^{1+\sigma}}.\end{aligned}$$

The last equivalence is true because there exist constants $C, c > 0$ independent of σ such that

$$\int_0^{1/|\sin x/2|}\frac{s^{2-\sigma}}{s^4+1}\,ds \le \int_0^1 s^{2-\sigma}\,ds + \int_1^\infty s^{-2-\sigma}\,ds \le C,$$

and

$$\int_0^{1/|\sin x/2|}\frac{s^{2-\sigma}}{s^4+1}\,ds \ge \frac{1}{2}\int_0^1 s^{2-\sigma}\,ds = c,$$

for all $1 \le \sigma < 2$. On the other hand, $\displaystyle\int_1^\infty D_t(x)\,\frac{dt}{t^{1+\sigma}} \le C\int_1^\infty \frac{dt}{t^{1+\sigma}} \le C$, for all $1 \le \sigma < 2$. Hence, (3.11) holds.

Now we derive the pointwise formula for $(-\Delta)^{\sigma/2}f(x)$. Suppose that $f\in C^\infty(\mathbb{T})$. We have $(P_t-\mathrm{I})^2f(x)=(P_t-\mathrm{I})(P_tf(x)-f(x))=P_{2t}f(x)-2P_tf(x)+f(x)$. Using (2.5) and (3.6),

$$\begin{aligned}(-\Delta)^{\sigma/2}f(x)&=\frac{1}{c_\sigma}\int_0^\infty\int_{\mathbb{T}}(f(x)-f(y))D_t(x-y)\,dy\,\frac{dt}{t^{1+\sigma}}\\&=\frac{1}{c_\sigma}\int_0^\infty\int_{-\pi}^{\pi}(f(x)-f(x-z))D_t(z)\,dz\,\frac{dt}{t^{1+\sigma}}\\&=\frac{1}{c_\sigma}\int_0^\infty\int_{-\pi}^{\pi}(f(x)-f(x-z)-f'(x)z)D_t(z)\,dz\,\frac{dt}{t^{1+\sigma}},\end{aligned}$$

where in the last identity we used that $zD_t(z)$ is an odd function. Since $f\in C^{1,\sigma+\varepsilon-1}(\mathbb{T})$, we have that $|f(x)-f(x-z)-f'(x)z|\le C\|f\|_{C^{1,\sigma+\varepsilon-1}(\mathbb{T})}|\sin\frac{z}{2}|^{\sigma+\varepsilon}$. This and (3.11) allow us to see that the double integral above is absolutely convergent. Therefore,

$$(-\Delta)^{\sigma/2}f(x)=\int_{\mathbb{T}}(f(x)-f(y)-f'(x)(x-y))K^{\sigma/2}(x-y)\,dy,\qquad f\in C^\infty(\mathbb{T}).$$

with $K^{\sigma/2}$ as in (3.10). Noticing that the approximation argument in the proof of Lemma 3.3 can be applied here, we get the identity above for any $f\in C^{1,\sigma+\varepsilon-1}(\mathbb{T})$, and the integral is absolutely convergent. For the principal value, note that $zK^{\sigma/2}(z)$ is an odd function. □

Let us compute the second pointwise limit in (1.1).

Proposition 3.7. *Let $f\in C^2(\mathbb{T})$. Then*

$$\lim_{\sigma\to2^-}(-\Delta)^{\sigma/2}f(x)=-f''(x),\qquad x\in\mathbb{T}.$$

Proof. Recall the pointwise formula of Theorem 3.6. Let $Rf(x,y):=f(x)-f(y)-f'(x)(x-y)$. Then, for any $1\le\sigma<2$,

$$\begin{aligned}(-\Delta)^{\sigma/2}f(x)&=\int_{\mathbb{T}}\Big(Rf(x,y)-2f''(x)\left|\sin\tfrac{x-y}{2}\right|^2\Big)K^{\sigma/2}(x-y)\,dy\\&\qquad+2f''(x)\int_{\mathbb{T}}\left|\sin\tfrac{x-y}{2}\right|^2K^{\sigma/2}(x-y)\,dy=:J_{1,\sigma}+2f''(x)J_{2,\sigma}.\end{aligned}$$

By using the orthogonality of the trigonometric system on the torus,

$$\int_{-\pi}^{\pi}\left|\sin\tfrac{z}{2}\right|^2P_t(z)\,dz=\int_{-\pi}^{\pi}\frac{\cos z-1}{2}\,P_t(z)\,dz=\frac{e^{-t}-1}{2}.$$

With this, (3.10) and Tonelli's theorem we compute $J_{2,\sigma}$:

$$\begin{aligned}J_{2,\sigma}&=\frac{1}{c_\sigma}\int_{-\pi}^{\pi}\left|\sin\tfrac{z}{2}\right|^2\int_0^\infty(2P_t(z)-P_{2t}(z))\,\frac{dt}{t^{1+\sigma}}\,dz\\&=\frac{1}{c_\sigma}\int_0^\infty\left(2\int_{-\pi}^{\pi}\left|\sin\tfrac{z}{2}\right|^2P_t(z)\,dz-\int_{-\pi}^{\pi}\left|\sin\tfrac{z}{2}\right|^2P_{2t}(z)\,dz\right)\frac{dt}{t^{1+\sigma}}\\&=-\frac{1}{2c_\sigma}\int_0^\infty\left(1-e^{-t}\right)^2\frac{dt}{t^{1+\sigma}}=-\frac{1}{2}.\end{aligned}$$

Hence $(-\Delta)^{\sigma/2}f(x)=J_{1,\sigma}-f''(x)$. It remains to show that $J_{1,\sigma}\to0$ as $\sigma\to2^-$. Let ϵ be any positive number. Since f'' is uniformly continuous on $\mathbb{T}$, there exists $\delta=\delta(\epsilon)>0$ such that $|f''(x)-f''(y)|<\epsilon$ for all $x,y\in\mathbb{T}$ with $\left|\sin\frac{x-y}{2}\right|<\delta$. Hence

$$\begin{aligned}|J_{1,\sigma}|&\le\frac{C\epsilon}{c_\sigma}\int_{\left|\sin\frac{x-y}{2}\right|<\delta}\left|\sin\tfrac{x-y}{2}\right|^{1-\sigma}dy+\frac{C}{c_\sigma\delta^{1+\sigma}}\int_{\delta\le\left|\sin\frac{x-y}{2}\right|\le1}dy\\&\le\frac{C\delta^{2-\sigma}}{c_\sigma(2-\sigma)}\cdot\epsilon+\frac{C}{c_\sigma\delta^{1+\sigma}}\longrightarrow C\epsilon,\quad\text{as }\sigma\to2^-,\end{aligned}$$

because of (2.6). Since ϵ was arbitrary, $J_{1,\sigma}\to0$ as $\sigma\to2^-$ and the proof is complete. □

Remark 3.8. When $f \in C^{2,\alpha}(\mathbb{T})$, $0 < \alpha \leq 1$, we also have $\lim_{\sigma\to 2^+}(-\Delta)^{\sigma/2}f(x) = -f''(x)$. Indeed, for $2 < \sigma < 3$ we can write $\sigma = 2 + \epsilon$ for some $\epsilon > 0$. Then, by Proposition 3.5,

$$\lim_{\sigma\to 2^+}(-\Delta)^{\sigma/2}f(x) = \lim_{\epsilon\to 0^+}(-\Delta)^{\epsilon/2}(-\Delta)f(x) = (-\Delta)f(x) - \frac{1}{2\pi}\int_{\mathbb{T}}(-\Delta)f(y)\,dy = -f''(x).$$

This and Proposition 3.7 yield the second limit in (1.1).

Remark 3.9. Under the hypothesis of Theorems 3.4 and 3.6, formula (2.5) holds. Indeed, just write down the kernels in in (3.3) and (3.9) and apply Fubini's theorem.

4. Fractional Laplacian on Hölder spaces

The following theorem is analogous to the results by Silvestre [17] for $(-\Delta_{\mathbb{R}^n})^{\sigma/2}$ and Theorem A in [21] for $(-\Delta_{\mathbb{R}^n} + |x|^2)^{\sigma/2}$. It explains how the operators $(-\Delta)^{\sigma/2}$ interact with the Hölder spaces $C^{k,\alpha}(\mathbb{T})$. Our proof relies on semigroup ideas and is much simpler and general than the proofs in [17, 21] that are based on the manipulation of the pointwise formulas.

Theorem 4.1 (Interaction with Hölder spaces)**.** *Let $\alpha \in (0,1]$ and $0 < \sigma < 2$.*

1. *Let $f \in C^{0,\alpha}(\mathbb{T})$ and $\sigma < \alpha$. Then $(-\Delta)^{\sigma/2}f \in C^{0,\alpha-\sigma}(\mathbb{T})$ and*
$$\|(-\Delta)^{\sigma/2}f\|_{C^{0,\alpha-\sigma}(\mathbb{T})} \leq C\|f\|_{C^{0,\alpha}(\mathbb{T})}.$$
2. *Let $f \in C^{1,\alpha}(\mathbb{T})$ and $\sigma < \alpha$. Then $(-\Delta)^{\sigma/2}f \in C^{1,\alpha-\sigma}(\mathbb{T})$ and*
$$\|(-\Delta)^{\sigma/2}f\|_{C^{1,\alpha-\sigma}(\mathbb{T})} \leq C\|f\|_{C^{1,\alpha}(\mathbb{T})}.$$
3. *Let $f \in C^{1,\alpha}(\mathbb{T})$ and $\sigma \geq \alpha$, with $\alpha - \sigma + 1 \neq 0$. Then $(-\Delta)^{\sigma/2}f \in C^{0,\alpha-\sigma+1}(\mathbb{T})$ and*
$$\|(-\Delta)^{\sigma/2}f\|_{C^{0,\alpha-\sigma+1}(\mathbb{T})} \leq C\|f\|_{C^{1,\alpha}(\mathbb{T})}.$$
4. *Let $f \in C^{k,\alpha}(\mathbb{T})$ and assume that $k + \alpha - \sigma$ is not an integer. Then $(-\Delta)^{\sigma/2}f \in C^{l,\beta}(\mathbb{T})$, where l is the integer part of $k + \alpha - \sigma$ and $\beta = k + \alpha - \sigma + l$.*

In Theorem 4.1, (4) is a consequence of (1)–(3) by iteration. As we mentioned, Theorem 4.1 could be proved by following the ideas of [17] or [21] and taking into account that the conjugate function is bounded on $C^{0,\alpha}(\mathbb{T})$, $0 < \alpha < 1$, see Zygmund [23, Chapter III, (13.29)]. This method essentially uses the pointwise formulas (3.3) and (3.9).

We present a unified and more general way to prove Theorem 4.1. It stands on the semigroup characterization of $C^{k,\alpha}(\mathbb{T})$ given in Proposition 4.3 below. The method has several advantages. First, we only need to use the semigroup formula for the fractional powers (2.5), so the idea could be extended to other fractional operators, see for instance [13]. Secondly, Zygmund's classes can be also considered. The Zygmund space Λ_* contains the Lipschitz space $C^{0,1}(\mathbb{T})$, see [23, Chapter II]. Taking into account Proposition 4.3 below, we readily see that Theorem 4.1 is a direct corollary of Theorem 4.4.

Definition 4.2. Let $\beta > 0$ and $k = [\beta] + 1$. We define

$$\Lambda_\beta(\mathbb{T}) := \left\{f \in C(\mathbb{T}) : \|t^k\partial_t^k P_t f(x)\|_{L^\infty(\mathbb{T})} \leq At^\beta\right\}. \tag{4.1}$$

If A_k is the least constant A appearing above, then the norm in $\Lambda_\beta(\mathbb{T})$ is $\|f\|_{\Lambda_\beta(\mathbb{T})} = \|f\|_{L^\infty(\mathbb{T})} + A_k$.

Next we relate the spaces $\Lambda_\beta(\mathbb{T})$ with the Hölder spaces $C^{k,\alpha}(\mathbb{T})$ and the Zygmund class Λ_*.

Proposition 4.3. *Let $\beta > 0$.*

(i) Let $f \in C(\mathbb{T})$ and $k, \ell > \beta$ be two integers. Then the two conditions

$$\|t^k\partial_t^k P_t f(x)\|_{L^\infty(\mathbb{T})} \leq A_k t^\beta, \quad \|t^\ell\partial_t^\ell P_t f(x)\|_{L^\infty(\mathbb{T})} \leq A_\ell t^\beta$$

are equivalent. The least constants A_k and A_ℓ that satisfy the inequalities above are comparable.

(ii) If $0 < \beta < 1$ then $\Lambda_\beta(\mathbb{T}) = C^{0,\beta}(\mathbb{T})$, with equivalent norms.

(iii) *We have* $\Lambda_1(\mathbb{T})=\Lambda_*$, *the Zygmund class defined as the set of continuous functions* f *on* $\mathbb{T}$ *such that* $|f(x+h)+f(x-h)-2f(x)|\le C|h|$, *for all* $x\in\mathbb{T}$ *and* $h\in\mathbb{R}$. *The quantity*

$$\|f\|_{\Lambda_*}=\|f\|_{L^\infty(\mathbb{T})}+\sup_{|h|>0}\frac{\|f(x+h)+f(x-h)-2f(x)\|_{L^\infty(\mathbb{T})}}{|h|},$$

is equivalent to $\|f\|_{\Lambda_1(\mathbb{T})}$. *Consequently,* $C^{0,1}(\mathbb{T})\subsetneq\Lambda_1(\mathbb{T})$ *and* $\|f\|_{\Lambda_1(\mathbb{T})}\le C\|f\|_{C^{0,1}(\mathbb{T})}$.

(iv) *If* $1<\beta<2$ *then* $f\in\Lambda_\beta(\mathbb{T})$ *if and only if* f *is differentiable and* $f'\in\Lambda_{\beta-1}(\mathbb{T})$. *Moreover,* $\|f\|_{\Lambda_\beta(\mathbb{T})}$ *is equivalent to* $\|f\|_{L^\infty(\mathbb{T})}+\|f'\|_{\Lambda_{\beta-1}(\mathbb{T})}$. *Similarly,* $\Lambda_2(\mathbb{T})=\{f:f'\in\Lambda_*\}$.

(v) *If* β *is not an integer, then* $\Lambda_\beta(\mathbb{T})=C^{[\beta],\beta-[\beta]}(\mathbb{T})$ *with equivalent norms. Similarly, for* $\beta=j\in\mathbb{N}$, *we have* $\Lambda_j(\mathbb{T})=\left\{f:f^{(j)}\in\Lambda_*\right\}$.

Proof. Note that *(v)* follows from *(ii)–(iv)* by iteration. Items *(i)–(iv)* of this proposition, in the case when $\mathbb{T}$ is replaced by $\mathbb{R}^n$, is contained in Stein [18, Chapter V, Section 4]. The proof there relies on appropriate estimates and cancelations of the Poisson kernel on $\mathbb{R}^n$, the semigroup property of $e^{-t\sqrt{-\Delta_{\mathbb{R}^n}}}$ and the equation $(\partial_{tt}+\Delta_{\mathbb{R}^n})(e^{-t\sqrt{-\Delta_{\mathbb{R}^n}}}f)(x)=0$. The same ingredients are present in the case of the torus and the arguments from Stein follow line by line here. We just mention the estimates for the Poisson kernel on $\mathbb{T}$:

$$\left|\partial_t^kP_t(x)\right|\le C_k\left|\sin\tfrac{x}{2}\right|^{-(k+1)},\quad \left|\partial_t^kP_t(x)\right|\le C_ke^{-t}(1-e^{-t})^{-(k+1)},$$

for any $x\in\mathbb{T}$, $t>0$, and $k\ge1$, see [8, Lemma 3]. The derivatives with respect to x satisfy the same bounds. ☐

Theorem 4.4. *Let* $\beta>0$ *and* $0<\sigma<2$ *with* $\sigma<\beta$. *If* $f\in\Lambda_\beta(\mathbb{T})$ *then* $(-\Delta)^{\sigma/2}f\in\Lambda_{\beta-\sigma}(\mathbb{T})$, *and*

$$\|(-\Delta)^{\sigma/2}f\|_{\Lambda_{\beta-\sigma}(\mathbb{T})}\le C\|f\|_{\Lambda_\beta(\mathbb{T})}.$$

Proof. Suppose first that $0<\sigma<1$. We have to prove that

$$\|t^k\partial_t^kP_t(-\Delta)^{\sigma/2}f(x)\|_{L^\infty(\mathbb{T})}\le C\|f\|_{\Lambda_\beta(\mathbb{T})}t^{\beta-\sigma},\quad k=[\beta-\sigma]+1.$$

As $0<\sigma<1$, by Remark 3.9 and Proposition 4.3,

$$\begin{aligned}(-\Delta)^{\sigma/2}f(x)&=\frac{1}{c_\sigma}\int_0^\infty\left(P_sf(x)-f(x)\right)\frac{ds}{s^{1+\sigma}}\\&=\frac{1}{c_\sigma}\left(J_1(x,t)+J_2(x,t)\right),\end{aligned}$$

where $J_1(x,t)$ denotes the part of the integral from 0 to t. By using the semigroup property, the hypothesis and Proposition 4.3*(i)*,

$$\begin{aligned}|t^k\partial_t^kP_tJ_1(x,t)|&=\left|t^k\partial_t^kP_t\int_0^t\int_0^s\partial_rP_rf(x)\,dr\,\frac{ds}{s^{1+\sigma}}\right|\\&\le t^k\int_0^t\int_0^s\left|\partial_w^{k+1}P_wf(x)\big|_{w=t+r}\right|\,dr\,\frac{ds}{s^{1+\sigma}}\\&\le t^k\|f\|_{\Lambda_\beta(\mathbb{T})}\int_0^t\int_0^s(t+r)^{\beta-k-1}dr\,\frac{ds}{s^{1+\sigma}}\\&=t^\beta\|f\|_{\Lambda_\beta(\mathbb{T})}\int_0^t\int_0^{s/t}(1+u)^{\beta-k-1}\,du\,\frac{ds}{s^{1+\sigma}}\\&\le Ct^\beta\|f\|_{\Lambda_\beta(\mathbb{T})}\int_0^t\frac{s}{t}\,\frac{ds}{s^{1+\sigma}}=C\|f\|_{\Lambda_\beta(\mathbb{T})}t^{\beta-\sigma}.\end{aligned}$$

On the other hand, by the semigroup property and Proposition 4.3*(i)*,

$$\begin{aligned}|t^k\partial_t^k P_t J_2(x,t)| &\le t^k\int_t^\infty \left|\partial_w^k P_w f(x)\big|_{w=t+s}\right|\frac{ds}{s^{1+\sigma}}+\int_t^\infty \left|t^k\partial_t^k P_t f(x)\right|\frac{ds}{s^{1+\sigma}}\\ &\le C\|f\|_{\Lambda_\beta(\mathbb{T})}\left(t^k\int_t^\infty (t+s)^{\beta-k}\frac{ds}{s^{1+\sigma}}+t^{\beta-\sigma}\right)\\ &\le C\|f\|_{\Lambda_\beta(\mathbb{T})}t^{\beta-\sigma}\left(\int_1^\infty (1+u)^{\beta-k}\frac{du}{u^{1+\sigma}}+1\right)=C\|f\|_{\Lambda_\beta(\mathbb{T})}t^{\beta-\sigma}.\end{aligned}$$

Hence, the conclusion follows for $0<\sigma<1$.

Consider now the situation $1\le\sigma<2$. We can write $(-\Delta)^{\sigma/2}f=(-\Delta)^{\sigma/2-1/2}(-\Delta)^{1/2}f=(-\Delta)^{\sigma/2-1/2}R\partial_x f$, where $R=\partial_x(-\Delta)^{-1/2}$ is the conjugate function operator on $\mathbb{T}$. Observe that, by Proposition 4.3*(iv)*, if $f\in\Lambda_\beta(\mathbb{T})$, $\beta>\sigma\ge1$, then $\partial_x f\in\Lambda_{\beta-1}(\mathbb{T})$. Therefore, the result follows from the classical Korn-Privalov-Zygmund theorem that says that the conjugate function is bounded on the spaces $\Lambda_\gamma(\mathbb{T})$, $\gamma>0$, see Zygmund [23, Chapter III, (13.29)], and also from the case just proved above ($0<\sigma/2-1/2<1$). □

Part 2. **Extension problem and Harnack's inequalities**

5. The general extension problem

Let L be a nonnegative, densely defined and self-adjoint operator on $L^2(\Omega,d\eta)=L^2(\Omega)$, where Ω is an open subset of $\mathbb{R}^n$, $n\ge1$, and $d\eta$ is a positive measure on Ω. There is a unique resolution of the identity E, supported on the spectrum of L, such that

$$\langle Lf,g\rangle=\int_0^\infty \lambda\,dE_{f,g}(\lambda),\quad f\in\operatorname{Dom}(L),\ g\in L^2(\Omega).$$

Here $dE_{f,g}(\lambda)$ is a regular Borel complex measure of bounded variation. Throughout this section we use the notation $\langle f,g\rangle=\int_\Omega f(x)g(x)\,d\eta(x)$. The heat-diffusion semigroup generated by L is given by

$$\langle e^{-tL}f,g\rangle=\int_0^\infty e^{-t\lambda}\,dE_{f,g}(\lambda),\quad f,g\in L^2(\Omega),\ t\ge0.$$

Fix any $\gamma>0$. The fractional operators L^γ are defined by

$$\langle L^\gamma f,g\rangle=\int_0^\infty \lambda^\gamma\,dE_{f,g}(\lambda),\quad f\in\operatorname{Dom}(L^\gamma),\ g\in L^2(\Omega),$$

with domain

$$\operatorname{Dom}(L^\gamma)=\left\{f\in L^2(\Omega):\int_0^\infty \lambda^{2\gamma}\,dE_{f,f}(\lambda)<\infty\right\}\supset\operatorname{Dom}(L^{[\gamma]}).$$

Proof of Theorem 1.1. As in [20], the equality (1.3) means that $u(\cdot,y)\in\operatorname{Dom}(L)$ for any $y>0$ and

$$\langle u(\cdot,y),g(\cdot)\rangle=\frac{y^{2\gamma}}{4^\gamma\Gamma(\gamma)}\int_0^\infty\langle e^{-tL}f,g\rangle e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+\gamma}},$$

for all $g\in L^2(\Omega)$. It is proved in [20, Theorem 1.1] that, when $0<\gamma<1$, the function u given in (1.3) and interpreted as above, is well defined and satisfies (1.2) and (1.4). If we consider $\gamma>1$, then, by following exactly the same arguments as in [20], it is easy to see that u as in (1.3) is well defined and verifies (1.2). It remains to prove (1.4). We proceed by induction on $[\gamma]$. As we have just said, (1.4) is valid for $[\gamma]=0$. Assume (1.4) for $j<\gamma<j+1$, $j\in\mathbb{N}$. Let us check (1.4) for $j+1<\gamma+1<j+2$.

Take $f \in \mathrm{Dom}(L^{\gamma+1})$ and $g \in L^2(\Omega)$. Then

$$\begin{aligned}
\langle (y^{-1}\partial_y)u(\cdot,y), g(\cdot)\rangle &= \frac{1}{4^{(\gamma+1)}\Gamma(\gamma+1)}\int_0^\infty \langle e^{-tL}f,g\rangle (y^{-1}\partial_y)\left(y^{2(\gamma+1)}e^{-\frac{y^2}{4t}}\right)\frac{dt}{t^{1+(\gamma+1)}} \\
&= \frac{y^{2\gamma}}{4^{\gamma+1}\Gamma(\gamma+1)}\int_0^\infty \langle e^{-tL}f,g\rangle e^{-\frac{y^2}{4t}}\left(\frac{2(\gamma+1)}{t^{2+\gamma}} - \frac{y^2}{2t^{3+\gamma}}\right)dt \\
&= \frac{-2y^{2\gamma}}{4^{\gamma+1}\Gamma(\gamma+1)}\int_0^\infty \langle e^{-tL}f,g\rangle \partial_t\big(e^{-\frac{y^2}{4t}}t^{-(1+\gamma)}\big)\,dt \\
&= \frac{2}{4\gamma}\cdot\frac{y^{2\gamma}}{4^{\gamma}\Gamma(\gamma)}\int_0^\infty \langle e^{-tL}Lf,g\rangle e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+\gamma}} =: \frac{1}{2\gamma}\langle v(\cdot,y),g(\cdot)\rangle.
\end{aligned}$$

Observe that v is a solution to (1.2) with initial data $v(x,0) = Lf(x)$. By the induction hypothesis,

$$\begin{aligned}
&\lim_{y\to 0^+}\langle y^{1-2((\gamma+1)-(j+1))}\partial_y\left((y^{-1}\partial_y)^{j+1}u(\cdot,y)\right), g(\cdot)\rangle \\
&\quad = \frac{1}{2\gamma}\lim_{y\to 0^+}\langle y^{1-2(\gamma-j)}\partial_y\left((y^{-1}\partial_y)^{j}v(\cdot,y)\right), g(\cdot)\rangle \\
&\quad = \frac{1}{2\gamma}\frac{4^{\gamma-j}\Gamma(\gamma-j)}{2(\gamma-j)\Gamma(-(\gamma-j))}\cdot\frac{1}{2^j(\gamma-j)(\gamma-j+1)\cdots(\gamma-1)}\langle L^\gamma(Lf),g\rangle \\
&\quad = \frac{4^{(\gamma+1)-(j+1)}\Gamma((\gamma+1)-(j+1))}{2((\gamma+1)-(j+1))\Gamma(-((\gamma+1)-(j+1)))} \\
&\qquad \times \frac{1}{2^{j+1}((\gamma+1)-(j+1))((\gamma+1)-(j+1)+1)\cdots((\gamma+1)-1)}\langle L^{\gamma+1}f,g\rangle,
\end{aligned}$$

which is (1.4). The theorem is proved. □

Remark 5.1. As done in [20], more properties of the general extension problem could be established, like Poisson formulas, fundamental solutions, Cauchy–Riemann equations, conjugate Poisson kernels and L^p estimates.

Remark 5.2. If we assume that $f \in \mathrm{Dom}(L^{[\gamma]+1})$, $\gamma > 0$, then

$$L^\gamma f = \frac{1}{d_\gamma}\int_0^\infty \left(e^{-tL} - \mathrm{I}\right)^{[\gamma]+1} f\,\frac{dt}{t^{1+\gamma}}, \quad \text{in } L^2(\Omega),$$

with $d_\gamma = \int_0^\infty \left(e^{-t}-1\right)^{[\gamma]+1}\frac{dt}{t^{1+\gamma}}$. Therefore, in the case that L has an explicit heat kernel, one could obtain parallel pointwise formulas for $L^\gamma f(x)$ like those of Theorems 3.4 and 3.6.

6. HARNACK'S INEQUALITIES FOR $(-\Delta)^{\sigma/2}$

In this section we apply the extension problem to prove interior and boundary Harnack's inequalities for $(-\Delta)^{\sigma/2}$, $0 < \sigma < 2$.

Let us see how the extension problem for the fractional Laplacian on the torus admits a classical solution. We show this by using the classical Fourier method.

Take $f \in \mathrm{Dom}(-\Delta)$. We first claim that a solution $u : \mathbb{T}\times[0,\infty) \to \mathbb{R}$ to the extension problem (1.2) for f can be written as

$$\begin{aligned}
u(x,y) &= \frac{y^\sigma}{4^{\sigma/2}\Gamma(\sigma/2)}\int_0^\infty e^{t\Delta}f(x)e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+\sigma/2}} \\
&= \frac{y^\sigma}{4^{\sigma/2}\Gamma(\sigma/2)}\sum_{n\in\mathbb{Z}}c_n(f)e^{inx}\int_0^\infty e^{-tn^2}e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+\sigma/2}}.
\end{aligned} \tag{6.1}$$

Indeed, to see (6.1), observe that $e^{t\Delta}f(x) = \sum_{n\in\mathbb{Z}} e^{-tn^2}c_n(f)e^{inx}$, where the series converges uniformly in x, because

$$\sum_{n\in\mathbb{Z}} e^{-tn^2}|c_n(f)| \le \|f\|_{L^2(\mathbb{T})}\left(\sum_{n\in\mathbb{Z}} e^{-2tn^2}\right)^{1/2} \le \frac{C\|f\|_{L^2(\mathbb{T})}}{(1-e^{-ct})^{1/2}}.$$

Since

$$\int_0^\infty \sum_{n\in\mathbb{Z}} \left| e^{-tn^2}c_n(f)e^{-\frac{y^2}{4t}}\right| \frac{dt}{t^{1+\sigma/2}} \le C_f \int_0^\infty \frac{e^{-\frac{y^2}{4t}}}{(1-e^{-ct})^{1/2}}\frac{dt}{t^{1+\sigma/2}} < \infty,$$

Fubini's theorem can be applied to obtain the second equality of (6.1).

Secondly, $u(\cdot,y)\in C^2(\mathbb{T})$, for every $y>0$. Indeed, for $h>0$,

$$\frac{u(x+h,y)-u(x,y)}{h} = \frac{y^\sigma}{4^{\sigma/2}\Gamma(\sigma/2)}\sum_{n\in\mathbb{Z}} c_n(f)\frac{e^{in(x+h)}-e^{inx}}{h}\int_0^\infty e^{-tn^2}e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+\sigma/2}}.$$

As

$$\sum_{n\in\mathbb{Z}} |c_n(f)| \int_0^\infty ne^{-tn^2}e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+\sigma/2}} \le \sum_{n\in\mathbb{Z}} |c_n(f)|\int_0^\infty e^{-\frac{tn^2}{2}}e^{-\frac{y^2}{4t}}\frac{dt}{t^{1+(\sigma+1)/2}} < \infty,$$

by dominated convergence, u is differentiable with respect to x and the derivative can be taken inside the series in (6.1). A similar argument shows that $u(\cdot,y)\in C^2(\mathbb{T})$.

Finally, let us see that

$$\|y^{1-\sigma}u_y(x,y)\|_{L^2(\mathbb{T})} \to C_{\sigma/2}\|(-\Delta)^{\sigma/2}f\|_{L^2(\mathbb{T})}, \quad \text{as } y\to 0^+. \tag{6.2}$$

To prove (6.2) we use (6.1), the cancelation

$$\int_0^\infty e^{-\frac{y^2}{4t}}\left(\sigma - \frac{y^2}{2t}\right)\frac{dt}{t^{1+\sigma/2}} = 0, \quad y>0,$$

and dominated convergence as follows:

$$\begin{aligned}
\|y^{1-\sigma}u_y(x,y)\|^2_{L^2(\mathbb{T})} &= \sum_{n\in\mathbb{Z}} |c_n(f)|^2\left(\frac{1}{4^{\sigma/2}\Gamma(\sigma/2)}\int_0^\infty e^{-tn^2}e^{-\frac{y^2}{4t}}\left[\sigma-\frac{y^2}{2t}\right]\frac{dt}{t^{1+\sigma/2}}\right)^2\\
&= \sum_{n\in\mathbb{Z}} |c_n(f)|^2\left(\frac{1}{4^{\sigma/2}\Gamma(\sigma/2)}\int_0^\infty (e^{-tn^2}-1)e^{-\frac{y^2}{4t}}\left[\sigma-\frac{y^2}{2t}\right]\frac{dt}{t^{1+\sigma/2}}\right)^2\\
&\longrightarrow \sum_{n\in\mathbb{Z}} |c_n(f)|^2\left(\frac{\sigma}{4^{\sigma/2}\Gamma(\sigma/2)}\int_0^\infty (e^{-tn^2}-1)\frac{dt}{t^{1+\sigma/2}}\right)^2\\
&= C^2_{\sigma/2}\sum_{n\in\mathbb{Z}} |n|^{2\sigma}|c_n(f)|^2 = C^2_{\sigma/2}\|(-\Delta)^{\sigma/2}f\|^2_{L^2(\mathbb{T})}.
\end{aligned}$$

We also note that if $f\ge 0$ then $u\ge 0$.

Theorem 6.1 (Interior Harnack's inequality)**.** *Fix $0<\sigma<2$ and an interval $I\subseteq\mathbb{T}$. There exists a positive constant C depending only on σ such that for any subinterval J compactly contained in I,*

$$\sup_J f \le C\inf_J f,$$

for all $f\in\operatorname{Dom}(-\Delta)$, $f\ge 0$ in $\mathbb{T}$, such that $(-\Delta)^{\sigma/2}f=0$ in $L^2(I)$. As a consequence, any f verifying these properties is a continuous function in J.

Proof. Set $\tilde{u}(x,y)=u(x,|y|)$, $x\in\mathbb{T}$, $y\in\mathbb{R}$, where u is as in (6.1). Let us verify that $\tilde{u}$ is a nonnegative weak solution of

$$\operatorname{div}(|y|^{1-\sigma}\nabla\tilde{u}) = 0, \quad \text{in } Q := I\times(-R,R),\ R>0. \tag{6.3}$$

Indeed, for any $\varphi \in C_c^\infty(I \times (-R,R))$ and $\delta > 0$, by applying the divergence theorem,

$$\begin{aligned}\int_Q |y|^{1-\sigma}\nabla\tilde{u}\cdot\nabla\varphi\,dx\,dy &= \int_{Q\cap\{|y|\geq\delta\}} \operatorname{div}(|y|^{1-\sigma}\varphi\nabla\tilde{u})\,dx\,dy + \int_{Q\cap\{|y|<\delta\}} |y|^{1-\sigma}\nabla\tilde{u}\cdot\nabla\varphi\,dx\,dy \\ &= \int_I \varphi(x,\delta)\delta^{1-\sigma}\tilde{u}_y(x,\delta)\,dx + \int_{I\times(-\delta,\delta)} |y|^{1-\sigma}\nabla\tilde{u}\cdot\nabla\varphi\,dx\,dy.\end{aligned}$$

The first term above is bounded by $\|\varphi\|_{L^\infty((-R,R);L^2(I))}\|\delta^{1-\sigma}\tilde{u}_y(x,\delta)\|_{L^2(I)}$, which tends to 0 as $\delta \to 0^+$ because of (6.2). As for the second term, we write $\nabla\tilde{u}\cdot\nabla\varphi = \partial_x\tilde{u}\partial_x\varphi + \partial_y\tilde{u}\partial_y\varphi$, so the integral splits into two parts $J_1 + J_2$. To deal with J_1, we see that, as $f \in \operatorname{Dom}(-\Delta)$, the derivative $\partial_x f \in L^2(\mathbb{T})$. Next we check that $\|\partial_x u(x,y)\|_{L^2(\mathbb{T})} \to \|\partial_x f\|_{L^2(\mathbb{T})}$, as $y \to 0^+$. This is proved by using (6.1), a change of variables and dominated convergence:

$$\begin{aligned}\|\partial_x u(x,y)\|^2_{L^2(\mathbb{T})} &= \sum_{n\in\mathbb{Z}} n^2|c_n(f)|^2\left(\frac{y^\sigma}{4^{\sigma/2}\Gamma(\sigma/2)}\int_0^\infty e^{-tn^2}e^{-\frac{y^2}{4t}}\,\frac{dt}{t^{1+\sigma/2}}\right)^2 \\ &= \sum_{n\in\mathbb{Z}} n^2|c_n(f)|^2\left(\frac{1}{\Gamma(\sigma/2)}\int_0^\infty e^{-\frac{y^2}{4s}n^2}e^{-s}\,\frac{ds}{s^{1-\sigma/2}}\right)^2 \\ &\longrightarrow \sum_{n\in\mathbb{Z}} n^2|c_n(f)|^2 = \|\partial_x f\|^2_{L^2(\mathbb{T})}, \quad \text{as } y \to 0^+.\end{aligned}$$

Thus there exists a constant $C(f)$ such that $\|\partial_x\tilde{u}(x,y)\|_{L^2(\mathbb{T})} < C(f)$ for all sufficiently small y. Hence, $|J_1| \leq C(f,\varphi)\delta^{2-\sigma} \to 0$, as $\delta \to 0$. For J_2, by (6.2), there exists C such that $\||y|^{1-\sigma}\tilde{u}_y(x,y)\|_{L^2(I)} \leq C$ for all sufficiently small y. Therefore,

$$|J_2| \leq \int_{-\delta}^{\delta} \||y|^{1-\sigma}\tilde{u}_y(x,y)\|_{L^2(I)}\|\partial_y\varphi\|_{L^2(I)}\,dy \leq C_\varphi\delta \to 0, \quad \delta \to 0.$$

Hence, $\tilde{u}$ is a nonnegative weak solution to (6.3) in $Q = I \times (-R,R)$. Equation 6.3 is a degenerate elliptic equation with A_2 weight $|y|^{1-\sigma}$. By applying Harnack's inequality of [10] to $\tilde{u}$, we get the conclusion for f and f is a continuous function in J. □

Remark 6.2. A natural question that arises is how to apply the extension problem to get interior Harnack's inequality for $(-\Delta)^{\sigma/2}$ with $\sigma > 2$. First, we must note that some extra hypothesis on f should be added. Indeed, Harnack's inequality for the biharmonic operator $(-\Delta_{\mathbb{R}^n})^2$ holds if we also know that $(-\Delta_{\mathbb{R}^n})f \geq 0$, the counterexample being $f(x) = x_1^2$ in $B_2(0)$, see [6]. Secondly, if $\sigma > 2$, the degeneracy weight in the extension equation $\operatorname{div}(y^{1-\sigma}\nabla u) = 0$ does not belong to any A_p class and, up to our knowledge, Harnack's inequality in this case is not known.

Theorem 6.3 (Boundary Harnack's inequality)**.** *Let $0 < \sigma < 2$ and $f_1, f_2 \in \operatorname{Dom}(-\Delta)$ be two nonnegative functions on $\mathbb{T}$. Suppose that $(-\Delta)^{\sigma/2}f_i = 0$ in $L^2(I)$, for some interval $I = (x_0,x_1) \subset \mathbb{T}$, and $f_i = 0$ on $(x_0-\varepsilon, x_0]$, for some small $\varepsilon > 0$, $i = 1,2$. There is a universal constant C such that*

$$\sup_{x\in(x_0,x_0+\frac{\varepsilon}{2})}\frac{f_1(x)}{f_2(x)} \leq C \inf_{x\in(x_0,x_0+\frac{\varepsilon}{2})}\frac{f_1(x)}{f_2(x)}.$$

Moreover, $\frac{f_1}{f_2}$ is α-Hölder continuous in $(x_0, x_0+\varepsilon/2)$, for some universal $0 < \alpha < 1$.

Proof. Without loss of generality we can assume that $x_0 = 0$, $x_1 = \pi$ and $\varepsilon = 3/2$. Let $\tilde{u}_i(x,y) = u_i(x,|y|)$, where u_i is the extension of f_i as in Theorem 1.1. As in the proof of Theorem 6.1, $\tilde{u}_i$ satisfies the degenerate elliptic equation $\operatorname{div}(|y|^{1-\sigma}\nabla\tilde{u}_i) = 0$ in the weak sense in $(0,\pi)\times\mathbb{R}$. Moreover, $\tilde{u}_i$ verifies the equation in the weak sense in $(\mathbb{T}\times\mathbb{R})\setminus\{(x,0) : -\pi \leq x \leq 0\}$ and $\tilde{u}_i(x,0) = f_i(x)$ for all $x \in (-\pi,0)$.


$y>0$
$f_i = 0$
$(-\Delta)^{\sigma/2} f_i = 0$
$-\pi$
$-3/2$
0
π
$\mathrm{div}(|y|^{1-\sigma}\nabla \tilde{u}_i) = 0$, with $\tilde{u}_i = 0$ on the dashed line

Consider the map Ψ from $\mathbb{R}^2 \setminus \{(x,0) : x \leq 0\}$ into $\mathbb{R}^2 \setminus \{(x,y) : x \leq 0\}$ given in polar coordinates by

$$(r\cos\theta, r\sin\theta) \longmapsto (\Psi_1(r,\theta), \Psi_2(r,\theta)) = (r\cos\tfrac{\theta}{2}, r\sin\tfrac{\theta}{2}), \quad r>0,\ \theta\in(-\pi,\pi).$$

Define $U_i(X,Y) = \tilde{u}_i(\Psi_1^{-1}(X,Y), \Psi_2^{-1}(X,Y))$, for all $(X,Y) \in \mathbb{R}^2 \setminus \{(x,y) : x \leq 0\}$. Observe that U_i is a continuous function, hence $U_i(0,Y) = 0$ continuously for all $-3/2 < Y < 3/2$. Moreover, by making the change of variables in the equation for $\tilde{u}_i$, it can be readily seen that U_i is a nonnegative weak solution of $\mathrm{div}\,(B(X,Y)\nabla_{X,Y} U_i) = 0$ in $(0,\pi)\times\mathbb{R}$, where $B(X,Y) = \left|\frac{2XY}{\sqrt{X^2+Y^2}}\right|^{1-\sigma}$.



Since B is an A_2 weight in $\mathbb{R}^2$, the equation for U_i above is a degenerate elliptic equation. By the boundary Harnack's inequality of [9],

$$\sup_{(0,3/4)\times(-3/4,3/4)} \frac{U_1}{U_2} \leq C \inf_{(0,3/4)\times(-3/4,3/4)} \frac{U_1}{U_2},$$

and U_1/U_2 is α-Hölder continuous. Going back to $\tilde{u}_1$ and $\tilde{u}_2$ and restricting them to $y=0$ the result follows. □

REFERENCES

[1] K. Bogdan, The boundary Harnack principle for the fractional Laplacian, *Studia Math.* **123** (1997), 43–80.
[2] L. Caffarelli, J.-M. Roquejoffre and O. Savin, Nonlocal minimal surfaces, *Comm. Pure Appl. Math.* **63** (2010), 1111–1144.

[3] L. Caffarelli, S. Salsa and L. Silvestre, Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian, *Invent. Math.* **171** (2008), 425–461.

[4] L. Caffarelli and L. Silvestre, An extension problem related to the fractional Laplacian, *Comm. Partial Differential Equations* **32** (2007), 1245–1260.

[5] L. Caffarelli and A. Vasseur, Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation, *Ann. of Math. (2)* **171** (2010), 1903–1930.

[6] G. Caristi and E. Mitidieri, Harnack inequality and applications to solutions of biharmonic equations, in: Partial Differential Equations and Functional Analysis, 1–26, Oper. Theory Adv. Appl. **168**, Birkhäuser, Basel, 2006.

[7] S.-Y. A. Chang and M. d. M. González, Fractional Laplacian in conformal geometry, *Adv. Math.* **226** (2011), 1410–1432.

[8] Ó. Ciaurri and L. Roncal, Littlewood–Paley–Stein g_k-functions for Fourier–Bessel expansions, *J. Funct. Anal.* **258** (2010), 2173–2204.

[9] E. B. Fabes, C. E. Kenig and D. Jerison, Boundary behavior of solutions to degenerate elliptic equations, in: Conference on Harmonic Analysis in Honor of Antoni Zygmund, Vol. I, II (Chicago, Ill., 1981), 577–589, Wadsworth Math. Ser., Wadsworth, Belmont, CA, 1983.

[10] E. B. Fabes, C. E. Kenig and R. Serapioni, The local regularity of solutions of degenerate elliptic equations, *Comm. Partial Differential Equations* **7** (1982), 77–116.

[11] C. E. Gutiérrez, Harnack's inequality for degenerate Schrödinger operators, *Trans. Amer. Math. Soc.* **312** (1989), 403–419.

[12] N. S. Landkof, *Foundations of Modern Potential Theory*, Die Grundlehren der mathematischen Wissenschaften, Band 180, Springer-Verlag, New York-Heidelberg, 1972.

[13] T. Ma, P. R. Stinga, J. L. Torrea and C. Zhang, Regularity properties of Schrödinger operators, *J. Math. Anal. Appl.* **388** (2012), 817–837.

[14] L. Schwartz, *Théorie des Distributions*, (French), Publications de l'Institut de Mathématique de l'Université de Strasbourg, No. IX-X, Hermann, Paris, 1966.

[15] M. F. Shlesinger, G. M. Zaslavsky and J. Klafter, Strange kinetics, *Nature* **363** (1993), 31–37.

[16] L. Silvestre, Eventual regularization for the slightly supercritical quasi-geostrophic equation, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27** (2010), 693–704.

[17] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, *Comm. Pure Appl. Math.* **60** (2007), 67–112.

[18] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series **30**, Princeton Univ. Press, Princeton, New Jersey, 1970.

[19] P. R. Stinga, Fractional powers of second order partial differential operators: extension problem and regularity theory. PhD thesis, Universidad Autónoma de Madrid, Madrid, 2010.

[20] P. R. Stinga and J. L. Torrea, Extension problem and Harnack's inequality for some fractional operators, *Comm. Partial Differential Equations* **35** (2010), 2092–2122.

[21] P. R. Stinga and J. L. Torrea, Regularity theory for the fractional harmonic oscillator, *J. Funct. Anal.* **260** (2011), 3097–3131.

[22] K. Yosida, *Functional Analysis*, reprint of the sixth (1980) edition, Classics in Mathematics, Springer-Verlag, Berlin, 1995.

[23] A. Zygmund, *Trigonometric Series*, Vol. I, II, Cambridge Univ. Press, Cambridge-New York-Melbourne, 1977.

Departamento de Matemáticas y Computación, Universidad de La Rioja, 26004 Logroño, Spain
E-mail address: `luz.roncal@unirioja.es`

Department of Mathematics, The University of Texas at Austin, 1 University Station, C1200, Austin, TX 78712-1202, United States of America
E-mail address: `stinga@math.utexas.edu`